











CEPA

Address

4-10 Stanley Street Bendigo Victoria 3550

Postal Address

Box 2002

Bendigo Mail Centre BENDIGO VIC 3554

Phone

(03) 5443 6072

Fax

(03) 5443 8014

Website

www.cepa.com.au

Email

info@cepa.com.au

Normal Trading Hours

Monday to Friday

8.30am - 5.30pm

Saturday 9.00am - Noon

All prices shown include GST & valid at time of publication

BARASTOC

Fibre-Beet MASH

CONDITIONING FIBE

A low sugar, high fibre conditioning feed that targets overall

condition including; weight gain,

topline, muscle tone and function, coat and hoof health. ^{20kg} \$**62**.20 NEW



BARASTOC

GOLDEN YOLK

FREE RANGE

Nutritionally balanced layer feed to support hen health

and egg production from the point of lay onwards.

\$**26**.50





BACK IN STOCK



Eraquell

Cereal-based pellets that look and smell like pony cubes. Broad spectrum wormer that controls internal equine parasites and all three species of bots. SPECIAL PRICE \$30.00 Until Sold Out



MEMBERS NOVEMBER SPECIAL

MecWormer plus Bots 100% worm kill (including benzimidazole resistant strains) SAVE \$32.5g \$3.00 SAVE \$32.5g \$19.50 CAUTION RESISTANCE ONSTITUENTS: 4 mpig ASANICTIA and 1875m ACTIVE CONSTITUENTS: 4 mpig ASANICTIA and 1875m ACTI



All Rounder

Balanced nutritional pellet diet suitable for horses and ponies being spelled or in light to moderate work.

\$**5**.00

^{20kg} \$**21**.50









NO DEET

NATURE'S

NATURE'S

Roll on

\$11.00

100g

Barrier

\$**17**.90

260g

Barrier \$**31**.00

> 125ml Spray

\$**20**.00

500ml

Spray \$40.50



Biosecurity Package Deal with Disinfectant

Includes: Bath, Mat, Disinfectant, Jug & Sign The Agboss Biosecurity Boot Bath package deals have been put together with the farmer in mind. The package includes a Boot Bath with Fingertip Rubber Mat, 5L Dual Action Disinfectant (makes 20 Litres) as well as a weatherproof Biosecurity Corflute Sign with eyelets.







HEALTHY HOOVES

Climate and weather vary depending on where in Australia you live. The hoof is a bit like a 'sponge', when it gets wet it increases in size and becomes softer and weaker. This means it is more fragile and if it stays wet it is more prone to cracks, breaks, infections and flares.

Aside from poor nutrition, lack of trimming and shoeing, excess moisture and genetics, horses that live in wet, humid environments for long periods of time are at the greatest risk. Hoof health depends on a balanced diet and a steady stream of nutrients.

Allow the hoof to dry out and have effective drainage in paddocks.



MYCOTOXINS

Certain fungi found on fresh pasture grasses, conserved roughage (hay, chaff, straw) and in grains can produce secondary compounds called mycotoxins. Mycotoxins favour environmental conditions, such as in warm and humid climates (as well as in improperly stored feeds). Once mycotoxins reach the hindgut they may affect the intestinal wall or enter the horse's blood stream, which can result in respiratory, gastrointestinal, neurological and reproductive issues. Contact CEPA for beneficial products.



SKIN FUNGUS

From the time of infection, ringworms can take up to 3 weeks for the first lesion to appear, and unfortunately during that time you may have already spread the condition onto other horses.

How is Ringworm transmitted?

By direct contact with Grooming Equipment, Sharing Tack, Infested Stables, Infested Floats and Rugs.

Prevention and Control?

Separate all the affected horses tack and disinfect and clean tack regularly with an effective disinfectant like Virkon S. Use separate grooming gear and separate the horse from any paddock mates until symptoms have gone. Wash your horse with an antifungal solution and use a gentle anti-fungal ointment.



STRINGHALT

Capeweed is prolific with the current conditions and produces high nitrate levels. Stringhalt is a known condition from eating too much capeweed or cat's ear. Horses show symptoms by lifting their legs and sometimes hitting their belly.

Horses should be moved from those paddocks but must not be confined! Lack of movement intensifies the condition. Encourage more magnesium in their diet with salt and calcium/magnesium.

Control the spread and spray affected areas with a broadleaf weed killer.







LIFE WISE











NOW AVAILABLE

